

Generating varieties for affine Grassmannians

Peter J. Littig and Stephen A. Mitchell

October 21, 2008

1 Introduction

Let G be a simply-connected compact Lie group with simple Lie algebra, and let $G_{\mathbb{C}}$ be its complexification. The affine Grassmannian associated to G is the homogeneous space $\mathcal{L}_G = \tilde{G}_{\mathbb{C}}/P$, where $\tilde{G}_{\mathbb{C}}$ is the affine Kac-Moody group of regular maps $\mathbb{C}^{\times} \rightarrow G_{\mathbb{C}}$ and P is the subgroup of maps $\mathbb{C} \rightarrow G_{\mathbb{C}}$. The space $\mathcal{L}_G$ is a projective ind-variety and has the homotopy type of ΩG , the space of based loops on G . Sitting inside $\mathcal{L}_G$ is a family of finite-dimensional, complex projective varieties, called *affine Schubert varieties*. These varieties are indexed by the elements of $\mathcal{Q}^{\vee}$, the coroot lattice associated to G . We write X_{λ} for the variety corresponding to $\lambda \in \mathcal{Q}^{\vee}$.

Its infinite-dimensionality notwithstanding, $\mathcal{L}_G$ is in many ways analogous to an ordinary Grassmannian. There are, however, two striking differences: (1) $\mathcal{L}_G$ is a topological group; and (2) the principal bundle defining $\mathcal{L}_G$ as a homogeneous space is topologically trivial. No finite-dimensional flag variety $G_{\mathbb{C}}/Q$ has these properties. These two properties of $\mathcal{L}_G$ give rise to interesting new phenomena that have no analogue in the classical setting. The most obvious is that there is a Schubert calculus not only for the cup product in cohomology, but also for the Pontrjagin product in homology. The problem is to compute the coefficients $a_{\sigma, \eta}^{\mu}$ in the expansion

$$[X_{\sigma}] \cdot [X_{\eta}] = \sum a_{\sigma, \eta}^{\mu} [X_{\mu}].$$

As far as we know, this problem was first considered in the second author's 1987 paper [18] on minuscule generating complexes, albeit in a very special case with specific topological applications in mind. Since then it has been studied by several authors, notably in unpublished work of Dale Peterson and in [13], [14], [15].

In the present paper we consider the Pontrjagin product on the point-set, geometric level. A finite-dimensional projective subvariety $X \subset \mathcal{L}_G$ is a *generating variety* if the image of $H_*X \rightarrow H_*\mathcal{L}_G$ generates $H_*\mathcal{L}_G$ as a ring. A Schubert variety X_{λ} is a *geometric generating variety* if (1) for all $n \geq 1$ the image of the n -fold multiplication $X_{\lambda}^n \rightarrow \mathcal{L}_G$ is a Schubert variety X_{λ_n} , with $\dim X_{\lambda_n} = n \cdot \dim X_{\lambda}$; and (2) $H_*X_{\lambda}^n \rightarrow H_*X_{\lambda_n}$ is surjective. Let α_0 denote the highest root of G .

Theorem 1.1 $X_{-\alpha_0^{\vee}}$ is a geometric generating variety, with $\lambda_n = -n\alpha_0^{\vee}$.

A classical theorem of Bott [3] can be reinterpreted to say that every $\mathcal{L}_G$ admits a smooth generating variety. In contrast, the generating varieties of the theorem above are all singular. Their advantage is that they are Schubert varieties, and give one canonical generating variety in each Lie type. It turns out that $\mathcal{L}_G$ admits a smooth Schubert generating variety if and only if G is not of type E_8 , F_4 , or G_2 (Proposition 6.6). In type C_n , it turns out that $X_{-\alpha_0^\vee}$ is essentially the generating complex for $\Omega Sp(n)$ considered by Hopkins [9].

The proof of Theorem 1.1 makes use of the balanced product $\tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G$. We construct an ind-variety structure on the balanced product, whose filtrations are topologically trivial algebraic fiber bundles $E_{\sigma,\lambda}$ with base X_σ and fiber X_λ (λ an anti-dominant element of the coroot lattice). Then we relate the group multiplication $\mathcal{L}_G \times \mathcal{L}_G \rightarrow \mathcal{L}_G$ to the action map $\tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G \rightarrow \mathcal{L}_G$.

Along the way we consider the surprising topological properties (1) and (2) and how they fail in the category of ind-varieties. In particular, we show that there is no topological group structure on $\mathcal{L}_G$ whose multiplication map is a morphism of ind-varieties (Corollary 3.4). Regarding property (2), although it is clear that the principal P -bundle $\tilde{G}_{\mathbb{C}} \rightarrow \tilde{G}_{\mathbb{C}}/P$ is not trivial as ind-varieties, it is also easy to show that the balanced product bundle is isomorphic to $\mathcal{L}_G \times \mathcal{L}_G$ as ind-varieties. It does not follow, however, that the bundles $E_{\sigma,\lambda}$ are products.

2 Notation and Conventions

Throughout this paper, G is a simple, simply-connected, compact Lie group with maximal torus T , Weyl group W , and complexification $G_{\mathbb{C}}$. We fix a Borel group B containing the Cartan subgroup $H = T_{\mathbb{C}}$, and let B^- denote the opposite Borel group. Let U be the unipotent radical of B , so that $B = HU$. Let $S \subset W$ denote the simple reflections, root system Φ , and simple roots $\alpha_s, s \in S$. Let $\mathcal{Q}^\vee$ denote the coroot lattice. We denote by $\mathcal{C}$ the dominant Weyl chamber determined by B .

Loop Groups. The primary references for this section are [21] and [12]; see also [19].

An *algebraic loop in $G_{\mathbb{C}}$* is a regular map $\mathbb{C}^\times \rightarrow G_{\mathbb{C}}$. The set of all such loops, which we denote by $\tilde{G}_{\mathbb{C}}$, is a topological group with respect to pointwise multiplication. In type A_{n-1} , for example, $G_{\mathbb{C}} = SL_n \mathbb{C}$ and $\tilde{G}_{\mathbb{C}} = SL_n(\mathbb{C}[z, z^{-1}])$. In general Lie type, $\tilde{G}_{\mathbb{C}}$ can be shown to be a complex affine ind-group.

Let $\Omega_{alg} G$ be the subgroup

$$\Omega_{alg} G = \{f \in \tilde{G}_{\mathbb{C}} : f|_{S^1} \in \Omega G\}.$$

We call $\Omega_{alg} G$ the group of based algebraic loops on G . Now, let P be the subgroup of $\tilde{G}_{\mathbb{C}}$ of algebraic loops $\mathbb{C} \rightarrow G_{\mathbb{C}}$ (i.e., algebraic loops that may be extended over the origin.) The following decomposition theorem is a fundamental part of the structure theory of loop groups.

Theorem 2.1 [*Iwasawa Decomposition*] *The multiplication map $\Omega_{alg} G \times P \rightarrow \tilde{G}_{\mathbb{C}}$ is a homeomorphism.*

We call this the Iwasawa decomposition because it is a modified analogue of Iwasawa's KAN decomposition for real Lie groups. The unmodified analogue takes the form $L_{alg} G \times$

$A \times \tilde{U} \xrightarrow{\cong} \tilde{G}_{\mathbb{C}}$, where A comes from the KAN decomposition of the constant loops $\tilde{G}_{\mathbb{C}}$, and $\tilde{U} = \{f \in \tilde{B} : f(0) \in U^-\}$.

Two corollaries of Theorem 2.1 are particularly important for our purposes. First, the homogeneous space $\mathcal{L}_G$ is homeomorphic to $\Omega_{alg}G$ and hence carries the structure of a topological group. As we shall observe later, such a group structure is wholly absent in the classical setting of homogeneous spaces of complex algebraic groups. Second, the principal bundle $P \longrightarrow \tilde{G}_{\mathbb{C}} \longrightarrow \mathcal{L}_G$ is trivial.

The following theorem, due, independently, to Quillen and Garland-Raghunathan, establishes the connection between $\Omega_{alg}G$ and the ordinary loop group ΩG .

Theorem 2.2 [Quillen, Garland-Raghunathan] *The inclusion $\Omega_{alg}G \longrightarrow \Omega G$ is a homotopy equivalence.*

Next, we describe the Tits system associated to $\tilde{G}_{\mathbb{C}}$. Let $\tilde{B} = \{f \in P : f(0) \in B^-\}$. The group $\tilde{B}$ is called the *Iwahori subgroup* of $\tilde{G}_{\mathbb{C}}$ and may be thought of as an affine analog of a Borel subgroup of $G_{\mathbb{C}}$. Let $\tilde{N}_{\mathbb{C}}$ denote the group of algebraic loops in $G_{\mathbb{C}}$ that take their image in $N_{\mathbb{C}}$. The intersection $\tilde{B} \cap \tilde{N}_{\mathbb{C}}$ is normal in $\tilde{N}_{\mathbb{C}}$ and the quotient group $\tilde{W} = \tilde{N}_{\mathbb{C}}/(\tilde{B} \cap \tilde{N}_{\mathbb{C}})$ is canonically identified with the affine Weyl group associated to G . Thus $\tilde{W}$ has a semi-direct product decomposition $\tilde{W} = \mathcal{Q}^{\vee} \rtimes W$. Here we note that since G is simply-connected, the lattice $\mathcal{Q}^{\vee} \subset \mathfrak{t}_{\mathbb{C}}$ can be identified with the subgroup $Hom(S^1, T) \subset \Omega_{alg}G$. Consequently, we alternate frequently between additive notation and multiplicative notation; the meaning will be clear from the context. It is shown in [19] that $(\tilde{G}_{\mathbb{C}}, \tilde{B}, \tilde{N}_{\mathbb{C}}, \tilde{S})$ is a topological Tits system.

The Affine Weyl Group. As a Coxeter system, the affine Weyl group has generating set $\tilde{S} = S \cup \{s_0\}$, where s_0 is reflection across the hyperplane $\alpha_0 = 1$. The cosets $\tilde{W}/W$ have two canonical sets of representatives: The coroot lattice $\mathcal{Q}^{\vee}$ and the minimal length elements $\tilde{W}^S$. An element λ of the coroot lattice is said to be *antidominant* if it lies in the closure of $-\mathcal{C}$, the *antidominant* Weyl chamber. We write $\mathcal{Q}_-^{\vee}$ for the set of antidominant elements in $\mathcal{Q}^{\vee}$. The antidominant elements of the coroot lattice are characterized by the following proposition.

Proposition 2.3 *For $\lambda \in \mathcal{Q}^{\vee}$, the following conditions are equivalent:*

- (a) $\lambda \in \tilde{W}^S$.
- (b) λW is maximal in $W \cdot \lambda W/W$.
- (c) λ is antidominant.

Proposition 2.4 *If $\sigma \in \tilde{W}^S$ and $\lambda \in \mathcal{Q}_-^{\vee}$, then*

$$l(\sigma\lambda) = l(\sigma) + l(\lambda).$$

The Affine Grassmannian. The affine Grassmannian associated to G is the homogeneous space $\mathcal{L}_G = \tilde{G}_{\mathbb{C}}/P$. By the Iwasawa decomposition, the natural map $\Omega_{alg}G \longrightarrow \mathcal{L}_G$ is a

homeomorphism, and hence $\mathcal{L}_G$ is homotopy equivalent to ΩG by the Quillen/Garland-Raghunathan theorem. As the notation suggests, $\tilde{G}_\mathbb{C}/P$ is closely analogous to a flag variety of a linear algebraic group. For example, the $\tilde{B}$ -orbits in $\mathcal{L}_G$ are finite-dimensional affine cells called Schubert cells. Each such cell is of the form $e_\sigma = \tilde{B} \cdot \sigma P / P$ where σ is an element of $\tilde{W}^S$ or $\mathcal{Q}^\vee$. More precisely, let $P^- \subset \tilde{G}_\mathbb{C}$ denote the subgroup of maps regular at infinity, let $P^{(-1)} = \{f \in P^- : f(\infty) = I\}$, and let $\tilde{U}_\sigma = \{u \in \tilde{U} : \sigma^{-1}u\sigma \in P^{(-1)}\}$. Then $\tilde{U}_\sigma$ is a finite dimensional unipotent group and hence is isomorphic to an affine space; furthermore, the natural map $\tilde{U}_\sigma \sigma \rightarrow e_\sigma$ is an isomorphism of varieties.

Given $\sigma \in \tilde{W}^S$, if n_σ is a lift of σ to $\tilde{N}_\mathbb{C}$, we shall follow tradition and write σP to denote the coset $n_\sigma P$. Also, we will write $\sigma P / P$ for the image of σP under the canonical projection $\tilde{G}_\mathbb{C} \rightarrow \mathcal{L}_G$.

The closure e_σ , which is denoted X_σ , is called an *affine Schubert variety* or simply, a Schubert variety. These varieties are (finite-dimensional) complex projective varieties.

3 $\mathcal{L}_G$ as a Topological Group

Since $\Omega_{alg} G$ is a topological group under pointwise multiplication, $\mathcal{L}_G$ inherits a topological group structure from the canonical homeomorphism $\Omega_{alg} G \xrightarrow{\cong} \mathcal{L}_G$. This group structure is a special feature of our present, affine setting. Indeed, a classical homogeneous space of the form $G_\mathbb{C}/P$ cannot even be given the structure of an H -space. A classical theorem of Hopf shows that the rational cohomology ring of a finite H -space is isomorphic to an exterior algebra on odd-dimensional classes. The cohomology of $G_\mathbb{C}/P$, however, is concentrated in even dimensions.

On the other hand, there is no algebraic ind-group structure on $\mathcal{L}_G$. Here *algebraic ind-group* means “group object in the category of ind-varieties”; it does not mean that the filtrations defining the ind-variety structure are subgroups. We say that an ind-variety X is irreducible if it cannot be written as a union of two proper Zariski closed subsets. It is easy to show that X is irreducible if and only if it admits a filtration by finite-dimensional irreducible subvarieties.

Proposition 3.1 *Let G be an irreducible projective algebraic ind-group. Then G is abelian.*

Proof: Let G_n be a filtration of G by irreducible projective varieties (note that G_n need not be a subgroup). Let $\gamma : G \times G \rightarrow G$ be the map that sends (g, h) to its commutator $ghg^{-1}h^{-1}$ and observe that γ is a morphism of ind-varieties. For $h \in G$, we let $\gamma_h(g) = \gamma(g, h)$ and note that $\gamma_e(g) = e$ for every $g \in G$. Since each G_n is irreducible and projective, Lemma 1 of [Shafarevich, 4.3] applies to the restriction of γ to $(G \times G)_n = G_n \times G_n$. Thus, γ_h is constant for each $h \in G_n$. Since $\gamma_h(e) = e$, it follows that each $\gamma_h \equiv e$, whence $\gamma(g, h) = e$ for all g, h in G_n . Since n may be chosen arbitrarily, we conclude that G is abelian.

Corollary 3.2 *The topological group structure on $\mathcal{L}_G$ is not an algebraic ind-group structure.*

We can say more, however.

Proposition 3.3 *Let G be an ind-variety with filtration (G_n) . Suppose that G admits a topological group structure with a regular multiplication map. If each G_n is a normal variety, then the inversion map is also regular.*

Proof: Let G satisfy the hypotheses stated in the proposition. Let $\mu : G \times G \rightarrow G$ be the multiplication map and $\chi : G \rightarrow G$ be the inversion map. For a fixed level G_p of the filtration, there exist integers q and n for which $\chi(G_p) \subset G_q$ and $\mu : G_p \times G_q \rightarrow G_n$ is a regular map of finite-dimensional varieties. Let $V \subset G_p \times G_q$ be the fiber of μ over the identity element e . Since μ is regular, V is a subvariety of $G_p \times G_q$. The projection $\pi_1 : V \rightarrow G_p$ is bijective and, since G_p is normal, an isomorphism ([12], p. 514). We conclude that $\chi = \pi_2 \pi_1^{-1} : G_p \rightarrow G_q$ is algebraic.

Corollary 3.4 *There does not exist a topological group structure on $\mathcal{L}_G$ with a regular multiplication map.*

Proof: Suppose that such a structure exists. Since the levels of the standard filtration on $\mathcal{L}_G$ are Schubert varieties—and thus normal ([12], p. 274)—Propositions 3.3 and 3.1 imply that the group structure is abelian. It follows that $\mathcal{L}_G$ is homotopy-equivalent to a product of Eilenberg-MacLane spaces ([8], 4K.7). Since the cohomology of $\mathcal{L}_G$ is torsion-free and concentrated in even degrees, the Eilenberg-MacLane spaces in question must be of type $K(\mathbb{Z}, 2)$. Thus $\mathcal{L}_G$ is a product of $\mathbb{C}P^\infty$'s. Since $\pi_2(\mathcal{L}_G) = \mathbb{Z}$, $\mathcal{L}_G \cong \mathbb{C}P^\infty$, a contradiction in any Lie type.

We next show that the inversion map on $\mathcal{L}_G$ is not regular. The proof makes use of the following lemma.

Lemma 3.5 *For $w \in W$, let $\varphi_w : G/T \rightarrow G/T$ denote the map $gT \mapsto gwT$. If $w \neq 1$, then φ_w is not a morphism of varieties $G_{\mathbb{C}}/B \rightarrow G_{\mathbb{C}}/B$.*

Proof: Suppose that $w \neq 1$ and that $\varphi_w : G_{\mathbb{C}}/B \rightarrow G_{\mathbb{C}}/B$ is regular. Let $r_w : G_{\mathbb{C}} \rightarrow G_{\mathbb{C}}$ denote right multiplication by w and consider the following diagram:

$$\begin{array}{ccc} G_{\mathbb{C}} & \xrightarrow{r_w} & G_{\mathbb{C}} \\ \pi \downarrow & & \downarrow \pi \\ G_{\mathbb{C}}/B & \xrightarrow{\varphi_w} & G_{\mathbb{C}}/B \end{array}$$

Observe that the diagram commutes when restricted to G in the upper left. Since G is Zariski dense in $G_{\mathbb{C}}$, the regularity of φ_w implies that the unrestricted diagram commutes. This yields a contradiction, however, since $\pi(U_w)$ is a point, whereas $\pi \circ r_w$ is an embedding on U_w .

Proposition 3.6 *The inversion map $\chi : \mathcal{L}_G \rightarrow \mathcal{L}_G$ is not regular.*

Proof: Note that, for $\lambda \in Q^\vee$, χ takes the P -orbit $\mathcal{O}_\lambda$ to the P -orbit $\mathcal{O}_{\lambda-\infty}$. Fix $\lambda \in Q_-^\vee$ with $\alpha_s \lambda = 0$ for all $s \in S$, and let Y_λ denote the $G_\mathbb{C}$ -orbit $G_\mathbb{C} \cdot \lambda$. The corresponding subset of $\Omega_{alg} G$ is Z_λ , the image of the map $\psi_\lambda : G/T \rightarrow \Omega_{alg} G$ given by $\psi_\lambda(gT) = g\lambda g^{-1}$. Since λ is dominant, the dominant representative of the orbit $\mathcal{O}_{\lambda-\infty}$ is $w_0 \cdot \lambda^{-1}$. We thus have a commutative diagram of homeomorphisms:

$$\begin{array}{ccc} G/T & \xrightarrow{\psi_\lambda} & Z_\lambda \\ \varphi_{w_0} \downarrow & & \downarrow \chi \\ G/T & \xrightarrow{\psi_{w_0 \cdot \lambda^{-1}}} & Z_\lambda \end{array}$$

Expressing this in terms of varieties yields

$$\begin{array}{ccc} G_\mathbb{C}/B & \longrightarrow & Y_\lambda \\ \varphi_{w_0} \downarrow & & \downarrow \chi \\ G_\mathbb{C}/B & \longrightarrow & Y_\lambda \end{array}$$

where now the horizontal maps are just the usual orbit maps $gB \mapsto g\lambda$, $gB \mapsto gw_0\lambda^{-1}$, respectively. If χ is algebraic in this diagram, then it is an isomorphism of varieties. This implies φ_{w_0} is algebraic, contradicting the lemma.

4 The balanced product $\tilde{G}_\mathbb{C} \times_P \mathcal{L}_G$

In this section we study the balanced product $\mathcal{E}_G = \tilde{G}_\mathbb{C} \times_P \mathcal{L}_G$. We begin with a discussion of the orbit map $\tilde{G}_\mathbb{C} \rightarrow \mathcal{L}_G$. Next we define a natural ind-variety structure on $\tilde{G}_\mathbb{C} \times_P \mathcal{L}_G$. Finally, we discuss the Pontrjagin product and the balanced product in homology.

4.1 The action map $\tilde{G}_\mathbb{C} \times \mathcal{L}_G \rightarrow \mathcal{L}_G$

The following proposition is a special case of [12], Lemma 7.4.10 and Proposition 7.4.11.

Proposition 4.1 *The orbit map $\pi : \tilde{G}_\mathbb{C} \rightarrow \mathcal{L}_G$ is a morphism of ind-varieties. Furthermore, π is a Zariski locally trivial principal P -bundle, as ind-varieties.*

Note that π is not globally trivial as ind-varieties, since $\tilde{G}_\mathbb{C}$ is affine and $\mathcal{L}_G$ is projective. On the other hand, it is a striking fact that π is globally trivial in the complex topology; this follows immediately from the Iwasawa decomposition. In contrast, for a finite-dimensional flag variety $G_\mathbb{C}/Q_I$ the principal Q_I bundle $G_\mathbb{C} \rightarrow G_\mathbb{C}/Q_I$ is never topologically trivial, since $\pi_2 G_\mathbb{C} \cong \pi_2 G = 0$, whereas $\pi_2(G_\mathbb{C}/Q_I) \cong H_2(G_\mathbb{C}/Q_I) \cong \mathbb{Z}^{|S|-|I|}$.

Call a morphism of ind-varieties $f : X \rightarrow Y$ a *quotient morphism* if whenever Z is an ind-variety and $g : Y \rightarrow Z$ is a set map such that gf is a morphism of ind-varieties, we have g a morphism of ind-varieties. Any regular map with local sections is a quotient morphism, hence:

Corollary 4.2 *π is a quotient morphism.*

This yields:

Proposition 4.3 *The action map $\theta : \tilde{G}_{\mathbb{C}} \times \mathcal{L}_G \rightarrow \mathcal{L}_G$ is a morphism of ind-varieties.*

The $\tilde{G}_{\mathbb{C}}$ -action on $\mathcal{L}_G$ has two nice properties: First, if $Q \subset \tilde{G}_{\mathbb{C}}$ is any proper, standard parabolic subgroup, then $\mathcal{L}_G$ is exhaustively filtered by Q -invariant Schubert varieties X_{λ} . Second, the action of Q on such an X_{λ} factors through a finite-dimensional quotient algebraic group Q/Q_{λ} . In other words, the action is “regular” in the sense of [12], p. 213, except that our parabolics are not pro-groups but rather subgroups of certain associated pro-groups.

In the case of interest to us, namely $Q = P$, there is a simple explicit description of the finite-dimensional quotients and the factored action. Thinking of P as $G(\mathbb{C}[z])$, there are natural surjective morphisms of algebraic ind-groups $\epsilon_k : P \rightarrow G[k]$, where $G[k] = G(\mathbb{C}[z]/z^k)$. Let $P^{(k)}$ denote the kernel of ϵ_k . Then if $\lambda \in Q_{-}^{\vee}$, it is not hard to show directly that $P^{(k)}$ acts trivially on X_{λ} for all sufficiently large k . Hence the finite-dimensional algebraic group $G[k]$ acts algebraically on X_{λ} .

We note also that $\tilde{G}_{\mathbb{C}}/P^{(k)} \rightarrow \mathcal{L}_G$ is a Zariski locally trivial principal $G[k]$ -bundle, as ind-varieties. Indeed for this it is enough to show that ϵ_k is locally trivial; but in fact it is globally trivial, as a principal $P^{(k)}$ -bundle of ind-varieties ([6]).

4.2 Ind-variety structure on $\mathcal{E}_G$

As topological spaces, we may filter $\tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G$ by the subbundles $\tilde{G}_{\mathbb{C}} \times_P X_{\lambda} \rightarrow \mathcal{L}_G$, where λ ranges over $Q_{-}^{\vee}$. Restricting these subbundles to X_{σ} , $\sigma \in \tilde{W}^S$, we obtain a local product $E_{\sigma, \lambda} \rightarrow X_{\sigma}$ with fiber X_{λ} . In fact $E_{\sigma, \lambda}$ has a unique structure of projective variety, such that $E_{\sigma, \lambda} \rightarrow X_{\sigma}$ is locally trivial in the Zariski topology as varieties. To prove this we need two lemmas.

Lemma 4.4 *Let $\pi : E \rightarrow X$ be a morphism of noetherian schemes over a noetherian base scheme A , and suppose that π is locally trivial with fiber F . Let $\mathcal{P}$ be any class of morphisms over A that is closed under composition and base change, and local in the target. Then if $F \rightarrow A$ and $X \rightarrow A$ are in $\mathcal{P}$, so is $E \rightarrow A$.*

In particular this holds when $\mathcal{P}$ is the class of separated, finite type, or proper morphisms.

Proof: The first part is an easy exercise. The last assertion follows from standard facts ([7], §II.4).

Lemma 4.5 *Let $\pi : E \rightarrow X$ be a local product with fiber F in the complex topology, and suppose that (i) X, F are varieties; (ii) there are local trivializations on a Zariski open cover U_{α} of X ; and (iii) the transition functions $U_{\alpha} \cap U_{\beta} \times F \rightarrow U_{\alpha} \cap U_{\beta} \times F$ associated to the*

cover in (ii) are regular. Then E has a unique algebraic variety structure compatible with the trivializations in (ii), so that π is a local product as varieties. Moreover, if X and F are complete, then so is E .

Proof: Identify the category of varieties over $\mathbb{C}$ with the category of reduced separated schemes of finite type over $\text{Spec } \mathbb{C}$. By the gluing lemma ([7], II.2.2) we obtain a reduced scheme E' over $\text{Spec } \mathbb{C}$, and a morphism of schemes $E' \rightarrow X$ that is locally trivial with fiber F . By the previous lemma, E' is separated of finite type, and is complete if X and F are complete. Furthermore, the set of complex points $E'(\mathbb{C})$ is naturally identified with E . This proves the lemma.

This yields a complete ind-variety structure on $\tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G$, indexed by the poset $\tilde{W}^S \times Q_-^{\vee}$. Note that the maps $\pi, \phi : \tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G \rightarrow \mathcal{L}_G$ given by $\pi([g, x]) = gP$, $\phi([g, x]) = gxP$ are morphisms of ind-varieties. Then $\Psi = (\pi, \phi)$ defines an isomorphism of ind-varieties

$$\Psi : \tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G \xrightarrow{\cong} \mathcal{L}_G \times \mathcal{L}_G.$$

The inverse map is given by $(gP, xP) \mapsto [gP, g^{-1}xP]$. Since $E_{\sigma, \lambda}$ is complete, it maps isomorphically onto its image in the projective ind-variety $\mathcal{L}_G \times \mathcal{L}_G$. Hence $E_{\sigma, \lambda}$ is projective, and $\tilde{G}_{\mathbb{C}} \times_P \mathcal{L}_G$ is a projective ind-variety.

Remarks: 1. We could have used Ψ to define the ind-variety structure, but the direct construction given above is more natural and has more general applicability. Note that Ψ embeds $E_{\sigma, \lambda}$ as a subvariety of $X_{\sigma} \times X_{\sigma, \lambda}$.

2. A similar argument shows that the $G[k]$ bundle $\tilde{G}_{\mathbb{C}}/P^{(k)} \rightarrow \mathcal{L}_G$ has a natural ind-variety structure, such that if $P^{(k)}$ acts trivially on X_{λ} , then $E_{\sigma, \lambda} \rightarrow X_{\lambda}$ is an algebraic fiber bundle with structure group $G[k]$.

4.3 Bi-Schubert cells in $E_{\sigma, \lambda}$

Fix $\sigma \in \tilde{W}^S$ and $\lambda \in Q_-^{\vee}$. Then $E_{\sigma, \lambda}$ is partitioned into *bi-Schubert cells*:

$$E_{\sigma, \lambda} = \coprod_{\tau \leq \sigma, \nu \leq \lambda} e_{\tau, \nu},$$

where $e_{\tau, \nu} = U_{\tau}\tau \times e_{\nu}$, identifying the latter with a subspace of $E_{\sigma, \lambda}$ in the evident way.

Lemma 4.6 a) Each bi-Schubert cell is a locally closed subvariety of $E_{\sigma, \lambda}$.

b) The closure relations on the bi-Schubert cells are given by the product order on $\tilde{W}^S \times \tilde{W}^S$.

Proof: The proof is similar to one of the standard proofs for the Schubert cells, so we only sketch the method. First we recall the Bott-Samelson varieties (see [12], §7.1). Let $\underline{s} = (s_1, \dots, s_p)$ be a sequence of elements of $\tilde{S}$, and let $Z_{\underline{s}}$ denote the balanced product

$$Z_{\underline{s}} = P_{s_1} \times_{\tilde{B}} P_{s_2} \times_{\tilde{B}} \dots P_{s_p} \times_{\tilde{B}} *,$$

where P_{s_i} denotes the parabolic subgroup generated by $\tilde{B}$ and s_i , and $*$ is a point. The Bott-Samelson variety $Z_{\underline{s}}$ is a smooth projective variety and an iterated $\mathbb{P}^1$ -bundle. We write $[g_1, \dots, g_p]$ for an equivalence class defining a point in $Z_{\underline{s}}$. Then there is a regular map $\rho : Z_{\underline{s}} \rightarrow \mathcal{L}_G$ given by $\rho[g_1, \dots, g_p] = g_1 g_2 \dots g_p P$. If $w = s_1 \dots s_p$ is an S -reduced expression, then ρ is a smooth resolution of X_w ; in fact it is an isomorphism over the top cell e_w .

Now suppose $\tau, \nu \in \tilde{W}^S$ as above. If $w = \tau\nu$ is a reduced product, choosing an S -reduced expression compatible with the product yields a Bott-Samelson map $\tilde{\rho} : Z_{\underline{s}} \rightarrow \bar{e}_{\tau, \nu}$ commuting in the diagram

$$\begin{array}{ccc} Z_{\underline{s}} & \xrightarrow{\rho} & X_w \\ \tilde{\rho} \downarrow & \nearrow \phi & \\ \bar{e}_{\tau, \nu} & & \end{array}$$

The proof is completed by analyzing $\tilde{\rho}$; details are left to the reader.

4.4 The balanced product in homology

We define the *balanced product* on $H_* \mathcal{L}_G$ by

$$[X_\tau] \star [X_\nu] = \phi_*[E_{\tau, \nu}].$$

Proposition 4.7 *The balanced product is associative but not commutative, and satisfies*

$$[X_\tau] \star [X_\nu] = \begin{cases} [X_{\tau\nu}] & \text{if } \tau\nu \text{ is } S\text{-reduced} \\ 0 & \text{otherwise} \end{cases}$$

Proof: Suppose $\tau\nu$ is reduced. Then since $E_{\tau, \nu} \rightarrow X_{\tau\nu}$ is an isomorphism of varieties over the top cell, it has degree one in homology. Hence $[X_\tau] \star [X_\nu] = [X_{\tau\nu}]$. If $\tau\nu$ is not reduced, we still have a Bott-Samelson map $\tilde{\rho} : Z \rightarrow E_{\tau, \nu}$, so that $\rho_*[Z] = [E_{\tau, \nu}]$. Then $\phi \circ \tilde{\rho} = \rho : Z \rightarrow \mathcal{L}_G$. Since the product is not reduced, it follows easily that $\dim \rho(Z) < \dim Z$ and hence $\rho_*[Z] = 0$. This proves the displayed formula. Clearly the product is associative, and is not commutative since $\tau\nu$ S -reduced need not imply $\nu\tau$ S -reduced.

Proposition 4.8 *Suppose $\sigma \in \tilde{W}^S$, $\lambda \in Q_-^\vee$. Then a) $\phi(E_{\sigma, \lambda}) = X_{\sigma\lambda}$*

b) $\phi_([E_{\sigma, \lambda}]) = [X_{\sigma\lambda}]$.*

c) $\phi_ : H_* E_{\sigma, \lambda} \rightarrow H_* X_{\sigma\lambda}$ is onto.*

d) For every $\omega \leq \sigma\lambda$, there exists $\tau \leq \sigma$ and $\nu \leq \lambda$ in $\tilde{W}^S$ such that $[X_\tau] \star [X_\nu] = [X_\omega]$.

Proof: Parts a, b and c follow easily from the cell decomposition and the Steinberg Lemma ([22], Theorem 15, and [11], Proposition 3.1). Part (d) is a refinement of (c), proved as follows: Suppose $\omega \leq \sigma\lambda$, and choose an S -reduced product decomposition $\omega = \tau\nu$ with $\tau \leq \sigma$ and $\nu \leq \lambda$. Then $\nu \in \tilde{W}^S$. We claim that if we choose such a decomposition with ν of maximal length, then also $\tau \in \tilde{W}^S$: For if not, then $\tau \downarrow \tau s$ for some $s \in S$, and hence there is a reduced product $\tau = \tau' s$. But then $\omega = \tau' s \nu$ is reduced, and $s\nu \leq s\lambda \leq \lambda$, since $\lambda \in Q_-^\vee$. This contradicts the maximality of ν , proving our claim. Thus $[X_\tau] \star [X_\nu] = [X_\omega]$.

4.5 The Pontrjagin product

Proposition 4.9 $E_{\sigma,\lambda} \rightarrow X_\sigma$ is topologically a product bundle $X_\sigma \times X_\lambda$. In fact there is a unique trivialization η such that the diagram

$$\begin{array}{ccc} X_\sigma \times X_\lambda & \xrightarrow{\eta} & E_{\sigma,\lambda} \\ \downarrow m & \searrow \phi & \\ \mathcal{L}_G & & \end{array}$$

commutes.

Proof: This follows from the Iwasawa decomposition.

Remark: These trivializations are not regular maps, since if they were then the product map m would be algebraic. We conjecture that $E_{\sigma,\lambda} \rightarrow X_\sigma$ is never a product bundle as varieties, nor indeed a product at all. We note also that η does not map $e_\sigma \times e_\lambda$ into $e_{\sigma,\lambda}$; we only have that $\eta(e_\sigma \times e_\lambda) \subset U_\sigma \sigma \times \mathcal{O}_\lambda$.

Proposition 4.10 Suppose $\sigma \in \tilde{W}^S$, $\lambda \in Q_-^\vee$. Then

- a) $m(X_\sigma \times X_\lambda) = X_{\sigma\lambda}$
- b) $m_*([X_\sigma] \otimes [X_\lambda]) = [X_{\sigma\lambda}]$.

Proof: Part (a) follows from Proposition 4.8 and Proposition 4.9. Since η is a map of X_λ -bundles over X_σ , having degree one on the base and the fiber, η itself has degree one. Part (b) then follows similarly.

5 The canonical generating variety

A *generating complex* for an H -space Y consists of a CW-complex K and a map $f : K \rightarrow Y$ such that the image of H_*f generates H_*Y as a ring. A *generating variety* for $\mathcal{L}_G$ is a finite-dimensional projective subvariety X such that $X \subset \mathcal{L}_G$ is a generating complex. A Schubert variety X_λ is a *geometric generating variety* if (1) for all $n \geq 1$ the image of the n -fold Pontrjagin multiplication $m^n : X_\lambda^n \rightarrow \mathcal{L}_G$ is a Schubert variety X_{λ_n} with $\dim X_{\lambda_n} = n \cdot \dim X_\lambda$; and (2) $m_*^n : H_*X_\lambda^n \rightarrow H_*X_{\lambda_n}$ is surjective. Since the sequence λ_n is cofinal in $\tilde{W}^S$, a geometric generating variety is in particular a Schubert generating variety. Now let $\lambda_0 = -\alpha_0^\vee \in Q_-^\vee$. The following theorem provides a canonical geometric generating variety in every Lie type.

Theorem 5.1 X_{λ_0} is a geometric generating variety for $\mathcal{L}_G$.

Proof: Take $\lambda_n = -n\alpha_0^\vee$. The theorem then follows from Proposition 4.10 and induction on n .

As a corollary we obtain a classical theorem of Bott [3].

Corollary 5.2 *The homology ring $H_*\Omega G$ is finitely-generated.*

Bott's proof uses generating complexes; indeed he introduced the concept partly for this purpose. His generating complexes, however, are smooth manifolds, whereas ours are always singular varieties as we will see shortly. We will also see that smooth Schubert generating varieties exist if and only if G is not of type E_8 , F_4 , or G_2 . Our canonical generating varieties have a number of interesting properties:

X_{λ_0} as Thom space: Taking P -orbits instead of $\tilde{B}$ -orbits in $\mathcal{L}_G$ yields a stratification

$$\mathcal{L}_G = \coprod_{\lambda \in \mathcal{Q}_-^\vee} \mathcal{O}_\lambda,$$

where $\mathcal{O}_\lambda$ is isomorphic to a vector bundle ξ_λ over the *Levi orbit* $M_\lambda = G_{\mathbb{C}} \cdot \lambda P$. Here M_λ is a flag variety $G_{\mathbb{C}}/Q_{I(\lambda)}$, where $I(\lambda) = \{s \in S : s\lambda W = \lambda W\}$. For details of this construction, see [20].

Taking $\lambda = \lambda_0$, one finds that $I(\lambda_0)$ corresponds to the nodes on the ordinary Dynkin diagram that are not adjacent to s_0 in the affine diagram. The adjoint action of $Q = Q_{I(\lambda_0)}$ on $\mathfrak{g}_{\mathbb{C}}$ stabilizes the one-dimensional root subalgebra $\mathfrak{u}_{\alpha_0}$, so that

$$\xi_{\lambda_0} = G_{\mathbb{C}} \times_Q \mathfrak{u}_{\alpha_0}.$$

Furthermore, since λ_0 is the minimal nonzero element of $\mathcal{Q}_-^\vee$, we have $X_{\lambda_0} = \overline{\mathcal{O}}_{\lambda_0} = \mathcal{O}_{\lambda_0} \cup *$, where $*$ is the basepoint. Hence X_{λ_0} is the one-point compactification of $\mathcal{O}_{\lambda_0}$ and so can be identified with the Thom space $T(\xi_{\lambda_0})$.

Minimal nilpotent orbits. Let $Y_0 \subset \mathfrak{g}_{\mathbb{C}}$ denote the minimal non-trivial nilpotent orbit; that is, the orbit of a nonzero element $x \in \mathfrak{u}_{\alpha_0}$. Then

$$Y_0 \cong G_{\mathbb{C}} \times_Q (\mathfrak{u}_{\alpha_0} - \{0\}).$$

In other words, Y_0 is the $\mathbb{C}^\times$ bundle associated to ξ_{λ_0} . Hence $X_{\lambda_0} - (M_{\lambda_0} \cup *) \cong Y_0$. Similarly, $\overline{Y}_0 \cong X_{\lambda_0} - M_{\lambda_0}$. In particular, $\overline{Y}_0$ and X_{λ_0} have equivalent singular points. Since the cone point in $\overline{Y}_0$ is known to be singular, this gives one way to see that X_{λ_0} is singular at the basepoint. We will give a different proof below. For an elegant computation of H^*Y_0 , see [10].

Birkhoff strata. The Birkhoff stratification of $\mathcal{L}_G$ is “dual” to the Schubert cells, and defines a descending filtration by ind-subvarieties Z_λ ($\lambda \in \mathcal{Q}^\vee \cong \tilde{W}^S$) [12] [21] [6]. There is a unique codimension one Birkhoff variety, namely $Z_{s_0} = Z_{\alpha_0^\vee}$. It is P -invariant and hence $Z_{\alpha_0^\vee} \cap X_{\lambda_0} \subset M_{\lambda_0}$; since both varieties are irreducible of codimension one, it follows that the inclusion is an equality. Thus if $\mathcal{U}_0$ is the Zariski open complement $\mathcal{L}_G - Z_{\alpha_0^\vee}$, we have $\mathcal{U}_0 \cap X_{\lambda_0} = \overline{Y}_0$. This observation enters into the beautiful classification of minimal degeneration singularities in [16].

Segments. Every $w \in \tilde{W}^S$ has a canonical reduced factorization $w = \sigma_1 \sigma_2 \dots \sigma_n$ into *segments*, defined in [2]. For present purposes we will use a slightly different definition, ignoring the

refinement for types BD of [2]. We say that $\sigma \in \tilde{W}^S$ is a *segment* if it is in the left W -orbit of s_0 in $\tilde{W}^S \cong \tilde{W}/W$. In other words, the segments are precisely the elements that index the cells of X_{λ_0} , excluding the basepoint. Each such σ has a unique reduced factorization $\sigma = \nu s_0$ with $\nu \in W^J$, where J is the complement in S of the set of nodes adjacent to s_0 in $\tilde{S}$. It is then easy to show that each $w \in \tilde{W}^S$ has a unique factorization into segments $w = \sigma_1 \sigma_2 \dots \sigma_n$ such that each partial product $\sigma_1 \dots \sigma_k$ is also in $\tilde{W}^S$. The canonical generating variety can be viewed as a geometric expression of this factorization; for example, it follows that there is a corresponding unique $\star$ -factorization of homology classes

$$[X_w] = [X_{\sigma_1}] \star [X_{\sigma_2}] \star \dots \star [X_{\sigma_n}].$$

No such factorization holds for the Pontrjagin product, however [13] [14].

Examples: (i) Type A_n , $n \geq 2$: This is the only type in which s_0 is adjacent to more than one node in the affine Dynkin diagram, and hence is the only type in which the flag variety M_{λ_0} is of non-maximal type. Explicitly, M_{λ_0} is the variety of flags of type $V^1 \subset V^n \subset \mathbb{C}^{n+1}$. Now let ξ_1, ξ_{n+1} denote the line bundles corresponding to V^1 and $\mathbb{C}^{n+1}/V^n$ respectively. Then $\xi_{\lambda_0} = \text{Hom}(\xi_{n+1}, \xi_1)$. In this case the canonical generating variety is not very efficient; the smallest generating varieties are the minuscule generating varieties $\mathbb{P}^n \subset \mathcal{L}_G$.

(ii) Type C_n , $n \geq 1$ (type A_1 is best thought of as type C_1 in this context). Here M_{λ_0} has type C_n/C_{n-1} ; thus $M_{\lambda_0} \cong \mathbb{P}^{2n-1}$. Then $\xi_{\lambda_0} \cong \eta^{*2}$ where η is the canonical line bundle (in topologist's but not geometer's terminology, i.e., η^* is the hyperplane section bundle). Hence $c_1(\xi_{\lambda_0}) \in H^2\mathbb{P}^{2n-1}$ is twice a generator. For later reference, we note that it follows that X_{λ_0} satisfies Poincaré duality rationally but not integrally.

Now let $\mu \in \mathcal{Q}^\vee$ be the unique element immediately below $\lambda_0 = -\alpha_0^\vee$; that is, $\mu = s_1 \lambda_0$. Since $H_* \Omega Sp(n) \cong \mathbb{Z}[a_1, \dots, a_n]$ with $|a_i| = 4i - 2$, it follows that X_μ is a Schubert generating variety for $\mathcal{L}_{Sp(n)}$. In fact $X_\mu = T(\xi_{\lambda_0} \downarrow \mathbb{C}P^{2n-2})$. This is the generating complex of [9].

(iii) Type G_2 . The flag variety M_{λ_0} is $G_{2\mathbb{C}}/Q$, where Q is the parabolic omitting the long node s_2 . It is a chain with cells indexed by $1, s_2, s_1 s_2, s_2 s_1 s_2, s_1 s_2 s_1 s_2, s_2 s_1 s_2 s_1 s_2$. For short, let X_k denote the Schubert variety of dimension k in $G_{\mathbb{C}}/Q_{s_0}$, and define $y_k \in H^{2k}(G_{\mathbb{C}}/Q_{s_0})$ by $\langle y_k, [X_k] \rangle = 1$. Then the Chevalley formula shows that $y_1 = c_1(\xi_{\lambda_0})$ and the cup products $y_1 y_{k-1} = a_k y_k$ with $a_k = 1, 3, 2, 3, 1$ for $1 \leq k \leq 5$. Proposition 6.2 below then shows that X_{λ_0} satisfies Poincaré duality rationally but not integrally, and in particular is not smooth.

6 Smooth vs. singular generating varieties

We first give an easy topological proof that X_{λ_0} is singular. From the Thom space description we see that X_{λ_0} is smooth away from the basepoint (i.e., the point at infinity).

Proposition 6.1 *a) X_{λ_0} never satisfies Poincaré duality over $\mathbb{Z}$. Hence the basepoint is a singular point.*

b) X_{λ_0} satisfies Poincaré duality over $\mathbb{Q}$ if and only if G has type A_1 , C_n , or G_2 .

This is a special case of [1], but it is worthwhile to sketch a proof here. Let X be a space satisfying Poincaré duality of dimension $2n$, with $H^{\text{odd}} X = 0$. We say that a space is

palindromic if it satisfies Poincaré duality additively, and is a *chain* if has Poincaré polynomial $1 + t^2 + t^4 + \dots + t^{2n}$. The proof of the following lemma is elementary, and left to the reader.


Lemma 6.2 *Let $\xi \downarrow X$ be a complex line bundle over X . Then*

- a) $T(\xi)$ is palindromic if and only if X is a chain if and only if $T(\xi)$ is a chain.*
- b) $T(\xi)$ satisfies Poincaré duality if and only if $H^*X \cong \mathbb{Z}[c_1\xi]/(c_1\xi)^{n+1}$.*

We also have:

Lemma 6.3 *$G_{\mathbb{C}}/Q_{I(\lambda_0)}$ is a chain if and only if G has type A_1 , C_n or G_2 .*

The proof is an easy and amusing exercise with Dynkin diagrams. Consider, for example, type E_8 :



The Weyl group elements indexing the cells of $G_{\mathbb{C}}/Q_{I(\lambda_0)}$ begin with s_8 , s_7s_8 , and so on until the fork at s_4 is reached and we find two elements of length 6. Hence $G_{\mathbb{C}}/Q_{I(\lambda_0)}$ is not a chain in type E_8 . Other types can be ruled out in the same way; in the case of F_4 note that the node s_3 counts as a fork since one can reverse direction along the doubled edge.

The two lemmas imply that if X_{λ_0} even rationally satisfies Poincaré duality, then G must have type A_1 , C_n , or G_2 . As shown in the previous section, these types do satisfy Poincaré duality rationally, but not integrally.

We now recall the classical construction of Bott [3], which yields certain smooth generating varieties for $\mathcal{L}_G$. These generating varieties are not always Schubert varieties, but they are always either Schubert varieties or Levi orbits. Bott works with the adjoint form G^{ad} of G , so that $Hom(S^1, T^{ad})$ can be identified with the coweight lattice $\mathcal{P}^{\vee}$. If $\lambda \in \mathcal{P}^{\vee}$, let C_{λ}^{ad} denote the centralizer of λ in G . The *Bott map*

$$B_{\lambda} : G^{ad}/C_{\lambda}^{ad} \longrightarrow \Omega_0 G^{ad}$$

is defined by $B_{\lambda}(gC_{\lambda}^{ad}) = \lambda g \lambda^{-1} g^{-1}$. Here Ω_0 denotes the basepoint component of the loop space. In fact Bott used the inverse of this map, but H_*B_{λ} and $H_*B_{\lambda}^{-1}$ generate the same subring of $H_*\Omega_0 G^{ad}$, so the distinction will not be important. Note that (i) the canonical map $\Omega G \longrightarrow \Omega_0 G^{ad}$ is a homeomorphism; and (ii) B_{λ} is in fact a map into the *algebraic* loops $\Omega_{alg,0} G^{ad}$.

The main result of [3] is that whenever λ is a “short minimal circle”, $G^{ad}/C_{\lambda}^{ad} \longrightarrow \Omega_0 G^{ad}$ is a generating complex. In particular, the fundamental coweight $\omega_s^{\vee}$ associated to a long simple root α_s is a short minimal circle; we will confine our attention to this special case. If $\lambda = \omega_s^{\vee}$ happens to lie in the coroot lattice $\mathcal{Q}^{\vee}$ (examples occur in every type except A , C),

then B_λ can be viewed in an evident way as a commutator map $G/C_\lambda \rightarrow \Omega_{alg} G$. Identifying $\Omega_{alg} G = L_{alg} G/G = \mathcal{L}_G$, we may also view B_λ as the map $gC_\lambda \mapsto \lambda g \lambda^{-1} P$. Since we are assuming $\lambda \in \mathcal{Q}^\vee$, multiplication by λ is homotopic to the identity and hence this last map is in turn homotopic to the map $gC_\lambda \mapsto g \lambda^{-1} P$. In other words, Bott's generating complex corresponds to the $G_{\mathbb{C}}$ -Levi orbit $M_{-\lambda}$. This yields the following corollary of Bott's work:

Proposition 6.4 *Suppose α_s is a long root and $\omega_s^\vee \in \mathcal{Q}^\vee$. Then the $G_{\mathbb{C}}$ -Levi orbit $M_{-\alpha_s^\vee}$ is a smooth generating variety for $\mathcal{L}_G$.*

In particular, if G is not of type A_n or C_n , then $M_{-\alpha_0^\vee}$ is a generating variety.

The second assertion follows because except in types AC , there is a unique node $t \in S$ adjacent to s_0 , linked to s_0 by a single edge. Hence α_t is long and $\omega_t^\vee = \alpha_0^\vee$.

At the opposite extreme, we can take s to be a minuscule node; that is, a node of S that is in the orbit of s_0 under the action of the automorphism group of the affine Dynkin diagram. (Warning: This is a possibly non-standard use of the term *minuscule*.) The fundamental coweights corresponding to the minuscule nodes are not in $\mathcal{Q}^\vee$, and indeed form a complete set of coset representatives for $\mathcal{P}^\vee/\mathcal{Q}^\vee$. Since these nodes always correspond to long roots, again Bott's construction yields generating varieties for $\Omega_0 G^{ad}$. The second author showed in [18] that these “minuscule generating complexes” correspond to smooth Schubert varieties that are in fact geometric generating varieties. In types E_8 , F_4 and G_2 , however, there are no minuscule nodes, so this construction does not apply. In view of the preceding proposition, we nevertheless obtain:

Proposition 6.5 *In all Lie types, $\mathcal{L}_G$ admits a smooth generating variety.*

Finally, the following proposition ties up a loose end from [18].

Proposition 6.6 *$\mathcal{L}_G$ admits a smooth Schubert generating variety if and only if G is not of type E_8 , F_4 or G_2 .*

Proof: What remains to be shown is that in the three exceptional types no smooth Schubert variety generates the homology. In fact we will show that no smooth Schubert variety can even generate the rational homology. Let $e_1 \leq e_2 \leq \dots \leq e_r$ denote the exponents of W , where $r = |S|$, and recall that $H_*(\Omega G; \mathbb{Q}) \cong \mathbb{Q}[a_1, \dots, a_r]$, where $|a_i| = 2e_i$. Hence if X_λ is a rational generating complex, we must have $\dim X_\lambda \geq e_r$. In the three exceptional types $e_r = 29, 11, 5$ respectively [4]. The smooth Schubert varieties were classified in [1]; in particular there are only finitely many in each Lie type. The maximal dimension d of a smooth Schubert variety X_λ is given by:

E_8 : $d = 14$ (X_λ is a quadric of type D_8/D_7)

F_4 : $d = 7$ (X_λ is a quadric of type B_4/B_3)

G_2 : $d = 2$ ($X_\lambda \cong \mathbb{P}^2$).

Hence none of these three types has a smooth Schubert generating variety.

References

- [1] Billey, S., and Mitchell, S., Smooth and palindromic Schubert varieties in affine Grassmannians, preprint 55pp: arXiv:0712.2871v1 (2007).
- [2] Billey, S., and Mitchell, S., Affine partitions and affine Grassmannians, preprint 46pp: arXiv:0712.2871 (2007).
- [3] Bott, R., The space of loops on a Lie group, Mich. Math. J. 5 (1958), 35-61.
- [4] Bourbaki, N., *Lie Groups and Lie Algebras*, Springer-Verlag, Berlin, 2002.
- [5] Evens, S. and Mirković, I., Characteristic cycles for the loop Grassmannian and nilpotent orbits, Duke Math. J. 97 (1999), 109-126.
- [6] Gutzwiller, L. and Mitchell, S.A., The topology of Birkhoff varieties, preprint 21pp: arXiv:0809.1091 (2008).
- [7] Hartshorne, R., *Algebraic Geometry*, Springer-Verlag, New York, 1977.
- [8] Hatcher, A., *Algebraic Topology*, Cambridge University Press, 2002.
- [9] Hopkins, M.J., Northwestern University Ph.D. thesis, 1984.
- [10] Juteau, D., Cohomology of the minimal nilpotent orbit, Transformation Groups 13 (2008), 355-387.
- [11] Kac, V., and Peterson, D., Defining relations of certain infinite-dimensional groups, in *Elie Cartan et les mathématiques d'aujourd'hui*, Astérisque, hors-série (1985), 165-208.
- [12] Kumar, S., *Kac-Moody Groups, their Flag Varieties and Representation Theory*, Birkhauser, Boston 2002.
- [13] Lam, T., Lapointe, L., Morse, J., and Shimozono, M., Affine insertion and Pieri rules for the affine Grassmannian, preprint: arXiv:math.CO/0609110v2 (2006).
- [14] Lam, T. Schilling, A., and Shimozono, M., Schubert polynomials for the affine Grassmannian of the symplectic group, preprint: arXiv:0710.2720v1 (2007).
- [15] Magyar, P., Schubert classes of a loop group, preprint: arXiv:0705.3826v1 (2007)
- [16] Malkin, A., Ostrik, V., and Vybornov, M., The minimal degeneration singularities in the affine Grassmannians, Duke Math. J. 126 (2005), 233-249.
- [17] Mitchell, S.A., A filtration of the loops on $SU(n)$ by Schubert varieties, Math Z. 193 (1986), 347-362.
- [18] Mitchell, S.A., The Bott filtration of a loop group, Proc. Barcelona Topology Conference, Lecture Notes in Math. v. 1298, 215-226, Springer-Verlag 1987.

- [19] Mitchell, S.A., Quillen's theorem on buildings and loop groups, *L'Enseignement Mathématique* 34 (1988), 123-166.
- [20] Mitchell, S.A., Parabolic orbits in flag varieties, preprint 2008.
- [21] Pressley, A., and Segal, G., *Loop Groups*, Clarendon Press, Oxford, 1986.
- [22] Steinberg, R., *Lectures on Chevalley Groups*, Yale University Lecture Notes, 1967.